



Purbeck Seabird Survey 2024

Sophie Lake & Emily Rush

FOOTPRINT ECOLOGY, FOREST OFFICE, BERE ROAD,
WAREHAM, DORSET BH20 7PA
WWW.FOOTPRINT-ECOLOGY.CO.UK
01929 552444



**FOOTPRINT
ECOLOGY**

Footprint Contract Reference: 807

Date: 20th August 2024

Version: Final

Recommended Citation: Lake, S. & Rush, E. (2024) Purbeck Seabird Survey 2024. Report by Footprint Ecology/National Trust.

Summary

This report presents data from the 2024 breeding seabird counts between Old Harry Rocks and St Aldhelm's Head. Counts have been carried out on the Purbeck coast since the mid-1960s and these data are presented in the context of both local trends over the last 50 years and national trends (up to 2019) for this stretch of the Dorset Coast. Poor weather meant that it was not possible to carry out the count between St Aldhelm's Head and White Nothe (usually carried out in May), so only the St Aldhelm's Head to Old Harry survey was conducted this year.

Seabirds breeding on the Purbeck coast include Fulmar, Cormorant, Shag, Herring Gull, Great Black-backed Gull, Kittiwake, Guillemot, Razorbill and Puffin. None of the populations are large. The Guillemot population remains the largest, with over 1,547 individuals counted on the breeding ledges in 2024, and the Puffin population is the smallest, with no birds observed during the survey (although up to 9 were reported on different occasions and 3 pairs may be breeding). Species such as Razorbill, Guillemot, and Puffin are thought to have been considerably more abundant in the first half of the 20th century, while Fulmar colonised, and Kittiwake increased markedly, during the second half of the 20th century, before declining. For a full discussion of previous Purbeck trends please see Lake et al. (2011).


Results from the 2024 monitoring show that:

- The Guillemot population increased on last year by 476 individuals to 1,547. This is similar to the peak count of 1,652 in 2022. All sub-populations increased on last year, including the re-emergence of a small population occupying White Ware, where no birds had been recorded since 1975.
- The Razorbill population likewise has increased compared to 2023, with 188 individuals counted on ledges, similar to that of the peak count in 2022. However, it is noticeable that sub-populations at Bird Rock and Funnel-Reform declined on the previous count and contrastingly Crab Hole and Durlston both saw an increase.
- The Herring Gull population remained stable in 2024 (an increase of one individual) following a continued decline in recent years. The population currently comprises 37% of the peak count since systematic recording began, and 12% of the highest count in 1969.
- The very small Great Black-backed Gull population increased by one this year (with an additional nest site found at Ballard Down), following the lowest level (since recording began in 2000) of just four apparently occupied nests the previous year.
- For Fulmar, the total number of adults on nests declined to levels previously seen in 2021. Nine individuals were recorded in Purbeck in 2024, with 11 individual nests recorded in total; the lowest level since recording began. Numbers are now at around 18% of the peak count in 2001, since systematic recording began. Breeding Fulmar are now present in just three of the six sites from where the species has regularly been recorded previously.

- The number of Cormorant nests at Ballard Down remains the same as in 2023, with nests noticeably scattered. This is the lowest number since the expansion of the 1980s (possibly attributable in part to the late survey date).
- Shag numbers remain stable (an increase of one) and are currently at the bottom end of recent fluctuations, and are at approximately 30% of the maximum count in 2006.
- Following a decrease in Kittiwake nests to their lowest count (4) in 2023, 5 nests were counted in 2024. This follows a slight increase between 2019 and 2021. The population has declined by 99% since a high point in 1982 and is in a precarious state.
- Although no Puffin were counted during the survey, observations from the cliff top suggest that their tiny population remains stable.

Challenging weather since 2021 has meant that counts of Cormorant nests at Gad Cliff have been difficult. Counts of individuals were carried out at Ballard to help provide a future estimate by looking at the ratio of birds to nests. There are however significant limitations to this approach due to fluctuating productivity between years and sites.

PURBECK SEABIRDS SURVEY 2024



Summary of breeding seabird population changes between Old Harry and St. Aldhelm's Head in Dorset. Counts are of apparently occupied nests/sites for all species except Guillemot, Razorbill and Puffin, which are of all individuals on breeding ledges. (Note that Fulmar and Shag counts before 2000 may not be complete).

Purbeck Seabirds Survey 2024

| | |
|---|-----------|
| Summary | ii |
| Purbeck Seabirds Survey 2024 | v |
| 1. Introduction..... | 1 |
| 2. Methods..... | 2 |
| 3. Results..... | 3 |
| Results of boat surveys | 3 |
| Estimate of number of breeding Puffins..... | 3 |
| 4. Discussion – comparison with previous years and UK trends | 6 |
| Fulmar | 7 |
| Cormorant..... | 9 |
| Shag..... | 11 |
| Herring Gull..... | 13 |
| Great Black-backed Gull..... | 15 |
| Kittiwake..... | 17 |
| Guillemot..... | 19 |
| Razorbill..... | 21 |
| Puffin..... | 24 |
| References | 26 |

Acknowledgements

This work builds on the remarkable legacy of seabird counts undertaken since the 1960s by the late Trev Haysom. 2024 survey work was carried out by Durwyn Liley, Phil Saunders, Emily Rush and Debbie Welham. An estimate of Puffin pairs was kindly provided by Richard Caldow. The work was funded by the National Trust and Durlston Country Park. Our thanks as always to Tom Greasty at Swanage Sea Fishing and the volunteer surveyors.

Cover photo © Durwyn Liley

1. Introduction

1.1 This report summarises the latest in a series of surveys (see Haysom, 1993; Haysom, 1977; Haysom, 1967; Lake et al. 2011; (Lake & Caals, 2022; Lake & Rush, 2023) of the breeding seabirds of the Dorset Coast. Surveys have been carried out annually, with the exception of 2020, when the survey was cancelled due to the Covid 19 pandemic. A count was nevertheless carried out between Durlston and St. Aldhelm's Head by Trev Haysom during this period (see Lake & Caals, 2022).

1.2 The stretch of coast is notable along the south coast of England in that it supports nine species of breeding seabird: Fulmar, Kittiwake, Cormorant, Shag, Great Black-backed Gull, Herring Gull, Guillemot, Razorbill, and Puffin. Eight of these species are Birds of Conservation Concern, the only exception being Cormorant. Kittiwake, Puffin, and Shag are now Red Listed, together with Herring Gull, while Guillemot, Razorbill, Great Black-backed Gull, and Fulmar are Amber Listed (Stanbury et al., 2021).

1.3 The South Dorset Coast is designated as a Site of Special Scientific Interest, Special Area of Conservation, and Jurassic Coast World Heritage Site for its wildlife and environmental interest, although breeding seabirds are not a designated feature.

2. Methods

2.1 It was possible to carry out both boat trips this year. The surveys took place on 27th May and 11th June 2024 solely between Old Harry and St. Aldhelm's Head, due to constraints related to the weather, choppy sea conditions and activity within the military Danger Area. Visibility was good on both counts, although the sea conditions were noticeably better for the second count.

2.2 Methods generally followed those recommended by Walsh *et al.* (1995). All observations of apparently occupied sites (AOS) or apparently occupied nests (AON) of Fulmar, Cormorant, Shag, Kittiwake, Herring Gull and Great Black-backed Gull were marked on enlarged photographs of the coast. Numbers of Guillemot, Razorbill and Puffin on known nesting ledges were counted and colonies were marked on enlarged photographs. The counts presented in the accompanying images and figures are from the June count when numbers were higher for the Guillemot and Razorbill.

2.3 The number of Puffins on the water and cliff ledges was also noted during the boat survey. However, the survey was undertaken during the day, when Puffin numbers tend to be at their lowest, as birds are either out at sea or out of sight within their breeding crevices. The number of breeding pairs was in the past been estimated by Trev Haysom using a consistent methodology (see Lake *et al.* 2011) – this was not possible in 2021 or 2022 but was undertaken by Richard Caldow in 2023 & 2024.

3. Results

Results of boat surveys

3.1 All AOS/AON and colonies are marked on the series of photographs supplied in the accompanying Photo Annex, whilst summary results are presented in the table at the start of this report. Survey sections follow those used historically and are given in (Lake et al., 2011).

Estimate of number of breeding Puffins

3.2 No Puffins were observed on the breeding ledge or water during the boat survey. The maximum number seen simultaneously by volunteer observers was eight on 2nd July and from weekly Birds of Poole Harbour seabirds boat trips (led by staff from Durlston Country Park) was nine individuals. Three breeding pairs were observed courting, mating and carrying nesting material, so it is probable that eggs were laid but that breeding was unsuccessful (R. Caldow, pers. comm).

Table 1: Breeding seabird records on the Dorset Coast, 2024. Counts are of apparently occupied nests or sites (AONs/AOSs) for all species except Guillemot and Razorbill (for which counts are of individuals at breeding sites) and Puffin (which is an estimate of pairs observed feeding chicks). Cormorant counts are for Ballard Down only due to difficulties observing nests at Gad Cliff and White Nothe – see discussion for further details.

| Species | 2024 total | Change since peak count | Change since last survey | Peak year | Peak count | Systematic surveys since: | Comments |
|-----------|------------|-------------------------|--------------------------|-----------|------------|---------------------------|---|
| Fulmar | 9 | -81.3% | -19 (-67.9%) | 2010 | 48 | 2001 | Colonised in 1940s, peaked in 1980s, despite a recent uptick in numbers, 2024 saw the lowest count of nests since recording began. |
| Cormorant | 15 | -91.3% | 0 (0%) | 1990 | 172 | 1964 | Declined to 1960s, sharp increase in 1990s followed steady decline to lowest count since 1981 increase. |
| Shag | 23 | -65.2% | 1 (4.6%) | 1992 | 66 | 1964 (partial) | Fairly steady despite fluctuations until 2011, subsequent apparent decline, particularly in the western sections. |
| Kittiwake | 5 | -98.4% | 1 (25%) | 1982 | 297 | 1957 | Rapidly increased throughout 1960s & 1970s, equally rapid decline, which slowed in the 2000s. 2023 count was the lowest since the expansion. |
| Guillemot | 1,547 | -6.4% | 476 (44.5%) | 2022 | 1,652 | 1964 | Large declines up to mid C20th, fluctuating then increasing notably from 2015 to a peak in 2022, remaining high in 2024 despite decrease in 2023. |

PURBECK SEABIRDS SURVEY 2024

| Species | 2024 total | Change since peak count | Change since last survey | Peak year | Peak count | Systematic surveys since: | Comments |
|-------------------------|------------|-------------------------|--------------------------|-----------|------------|---------------------------|---|
| Razorbill | 188 | -3.1% | 33 (21.3%) | 2022 | 194 | 1964 | Large declines up to mid C20th, sharp increase from 2008 to current high point (despite downturn in 2023). |
| Puffin ¹ | 3 | -100% | 0 (0%) | 1969 | 37 | 1967 | Large declines up to mid C20th which stabilised at current level around 1990. |
| Herring gull | 50 | -61% | 2 (4.2%) | 2015 | 128 | 2001 | Considerable decline 1960s - 1980s to low point in 2017, subsequent slight increase then decline to current stabilised levels in 2024. |
| Great black-backed gull | 4 | -77.8% | 4 (0%) | 2006 | 18 | 2001 | Fluctuating decline since 2001 with slight upturn of one AON per year in 2018 and 2019 followed by decline to lowest count recorded in 2023 and 2024. |

¹ Count from land supplied independently of boat survey.

4. Discussion – comparison with previous years and UK trends

4.1 Data from 1965 onwards were compiled and discussed in Lake *et al.* 2011. Here we update the dataset with the results of the 2024 survey between Old Harry and White Nothe.

4.2 The UK indices of abundance (JNCC, 2021)² show the relative change in population size, assigning a score of 100 to the population at the start date of the monitoring. These data were extracted from the [Seabird Monitoring Programme \(SMP\) Database](#)³. Data and have been provided to the SMP by the generous contributions of nature conservation and research organisations, and many volunteers throughout Britain and Ireland. Further information can be found in the individual UK trend chapters of the [online report](#)⁴. These indices are used to compare with a similar Dorset-based index of change for each species, to examine whether local trends differ from the national picture.

4.3 Contextual information on UK declines has been retained for readers who have not seen reports from previous years, but are indicated by the use of *grey italics*, enabling readers familiar with the text to skip information repeated between years. Please refer to Lake *et al.* 2011 for further context on each species and more information on historical records (including data constraints).

² The UK indices of abundance (JNCC, 2011) are compiled as part of the JNCC seabirds monitoring programme and earlier surveys in 1969-70 (Operation Seafarer), 1985-88 (Seabird Colony Register) and 1998-2002 (Seabird 2000).

³ <https://app.bto.org/seabirds>

⁴ <https://jncc.gov.uk/our-work/smp-report-1986-2019/>

Fulmar

After colonising Dorset in the 1940s, the number of breeding Fulmar increased to a peak in the 1980s. Since then, numbers have declined overall, despite short-term increases, similar to those seen in the previous two years. As such, the number of breeding Fulmer recorded in 2024 is the lowest count since recording began and just three of the six areas regularly used in the past were used in 2024. The decline in abundance broadly reflects that across the UK, although with wider fluctuations and a greater decline overall.

4.4 Fulmars breed between Ballard Down and St. Aldhelm's Head. Following the first record of Fulmars breeding on the Purbeck coast in 1943 (Haysom, 1977), numbers increased to a peak in the early 1980s. Comparable data are available from 2001, when the first complete surveys were undertaken (previous surveys did not include Ballard Down and counts prior to 1985 were of individuals rather than AONs). Since 2001, the overall trend appears to have been a decline, with peaks and troughs from year to year including a notable low point in 2021. Despite increase in 2022 and 2023, the Fulmer population observed in 2024 is the lowest count since systematic recording began, and matches numbers previously seen in 2021.




Figure 1: Changes in the numbers of apparently occupied breeding sites (AOSs) for Fulmar together with Purbeck and UK indices of abundance (counts before 1985 were of individuals rather than AOS). Counts before 2001 were only between Durlston and St Aldhelm's Head and have not been included in the Purbeck trend or minimum/maximum counts given in the summary table (Ballard Down has held between 2 and 23 nests).

4.5 Considering the available data (up to 2019), the local trend is similar to that of the UK as a whole (see Figure 1), although more variable due to the low numbers. A spectacular increase in the number and distribution of Fulmars in the UK and north Atlantic throughout the 20th century (prior to the mid-18th century, they bred in only one or two colonies in Iceland and in St Kilda) ceased in the last 20 years, and numbers then declined. The decline in Purbeck has been steeper.

4.6 The increase in Fulmar numbers in Europe is thought to have been driven by changes in food availability due to changes in temperature in the seas and to commercial fisheries, and to a reduction in human predation (Thompson, 2004). Subsequent declines in the UK have been attributed to changes in the North Sea whitefish industry, resulting in a decrease in

offal; and declines in sand eel populations in the North Sea and zooplankton in the Atlantic, likely due to climate change. Large numbers are also caught and accidentally killed by long-line fishing in the Norwegian Sea and North Atlantic. The Fulmar is Amber Listed due to the decline and degree of localisation of the breeding population.

Cormorant

The Handfast Point - Ballard Down population Cormorant population declined from a peak of 172 in 1990 to less than 10% of this figure in 2024, the lowest count recorded since the expansion in the 1980s. However, the low count may have been in part due to the relatively late survey date, by which time some nests may have disintegrated.

- 4.7 The number of Cormorants on the Dorset coast declined in the 1960's before increasing markedly in the 1980's. AONs at Ballard Down leapt from 10 in 1976 to 82 in 1982 and peaked at 172 in 1990. Since then, it has declined steadily, and numbers are now at the lowest recorded since the population expansion began, at just 15 AONs for 2023 and 2024 despite a small increase in 2022. However, the Cormorant count is usually taken from the May survey, which could not be carried out fully in 2024 – by 11th June some nests may have already disintegrated, and juveniles spread out meaning that the count may be an underestimate.
- 4.8 Nest locations typically vary between years, although a number of specific areas are often used on a rotational basis. In 2024, nest locations were notably scattered, rather than concentrated on the usually favoured sites.
- 4.9 Difficulties with visibility at Gad Cliff in previous years made it hard to count AONs. Therefore, the number of individuals on ledges were counted in order to provide an approximate index of change over time. Although a survey at Gad Cliff was not possible in 2024, individuals at Ballard Down were counted and totalled 35, including 17 chicks.




Figure 2: Cormorant AONs between Handfast Point and Ballard Down and the English indices of abundance (coastal populations only) and Purbeck indices of abundance.

4.10 *Because of significant regional variation in the abundance index (declines are particularly severe in northern Scotland), Figure 2 shows the trend for the Purbeck population for the years in which these data are available compared to the English index of abundance for coastal Cormorants. The Purbeck index decreased while the English index was still increasing, and the Purbeck population has also decreased further. The upturn in numbers nationally after 2011 was reflected in Purbeck in 2014-5, but Purbeck numbers then dropped again in contrast to the national trend, which appears more or less stable at the moment.*

4.11 *Nationally, increases in abundance up to 1995 are likely to have been facilitated by increased legal protection instigated under the Wildlife and Countryside Act 1981. Factors responsible for recent declines are likely to include increased mortality from licensed and unlicensed shooting, as well as possible changes in food availability (JNCC, 2011). Poor weather during the breeding season in 2012 and early in the breeding season in 2013 may have impacted on the Purbeck population, particularly at Ballard Down.*

Shag

The number of breeding Shags in Dorset is thought to have increased significantly from the early 20th century until the 1970s. Between the 1970s and 2010 the population remained fairly stable, although there were significant annual fluctuations. A clear decline until 2014 was followed by a stable but fluctuating count at a low level. The 2024 count remains low, with no nests recorded between Anvil Point and Ragged Rocks, and only one recorded at Blackers Hole (same as 2023). UK trends indicate a long-term decline.

4.12 Breeding Shags are generally scattered along the Purbeck coast. Records indicate an increase to a high point of 66 AONs in 2006. Since, then, more regular recording has shown a rapid decline to a low point of 21 birds in 2014. Numbers subsequently fluctuated between 21 and 35 nests and are now at 23 (see Figure 3).

4.13 No nests were recorded between Anvil Point and Ragged Rocks. However, a nest was once again recorded at Blackers Hole (where no nests were recorded for the first time in 2022). There was a notable shift in distribution of nests between Seacombe and Buttery Corner, where the numbers of AON's fluctuated compared to 2023.




Figure 3: Changes in the numbers of AONs for Shag together with local and UK indices of abundance.

4.14 The change in numbers of nesting Shags in Dorset has not closely reflected national trends, but both show an overall decline with fluctuations. The tendency for adults not to breed every year may be one reason for this variability. The Shag is Red Listed due to declines in the breeding population, and the international importance of both breeding and non-breeding populations in the UK (M. Eaton et al., 2015).

4.15 *In the UK overall, the Shag population increased slightly from the late 1960s to the mid-1980s (possibly due to increased legal protection, e.g. under the Wildlife and Countryside Act 1981, and reduced persecution (JNCC 2011)). It then gradually decreased, with an abrupt crash in 1994 and again in 2005 due to a wreck (mass mortality event) caused by food scarcity during a period of prolonged onshore gales on the east coast (Harris & Wanless, 1996). Note the initial steep rise in the index up to 1987 shown in Figure 3 is due to many adults choosing not to breed in 1986, resulting in low numbers at colonies that years.*

Herring Gull

Since systematic surveys began in 2000, the Herring Gull population has fluctuated, with a particularly steep decline since 2015. Although 2022 saw a notable increase, with the return of nesting birds to stretches of the coast where they were absent in 2021, 2024 remained at a low level similar to that of 2023 (the lowest count since 2000). Overall, there is thought to have been a marked decline in the Herring Gull population in Dorset in the second half of the twentieth century, which appears to be steeper than the national decline.

4.16 Records for Old Harry to St. Aldhelm's Head are only available from 2000 (see Figure 4). The patchy records available for Purbeck before this date suggest a decline (67% between 1965 and 1989) that is considerably more severe than the national decline (43% between the late-1960's and mid-1980's).

4.17 More systematic monitoring was introduced in 2000, by which time the population had recovered a little. However, a slow decline ensued, mirroring the overall UK trend (see Figure 4) until 2012 when numbers started increasing. In 2015, 128 nests were recorded, the most since 2000 (note that the total Dorset count in 2015 was still only around half of the number recorded in 1969). Numbers then reduced rapidly to a low of 55 AONs in 2021 and 50 in 2024, the second lowest level recorded since systematic recording began in 2001 (or any recording).




Figure 4: Number of AONs of Herring Gull and the Dorset and UK indices of abundance (UK monitoring started in 1986 and the UK index is based on coastal populations only).

4.18 *The Herring Gull is Red Listed in the UK due to a long-term decline in the population (Eaton et al., 2009). There has been a long-term decline in the coastal, natural-nesting, population within the UK. Factors implicated in the decline are botulism (thought to have been a major factor in the decline in the 1970's and 1980's), a decrease in the availability of food scavenged from refuse tips, and reductions in the availability of discards from fishing vessels, while ground predators have had an effect at some colonies.*

Great Black-backed Gull

The tiny Great Black-backed Gull population had remained fairly steady between 2000 and 2011, with numbers generally fluctuating between about 10 and 18, before dipping to fluctuate between 5 and 12 until 2023, when the population reached its lowest level of 4 nests since recording began. In 2024, 5 nests were recorded.

- 4.19 The numbers of Great Black-backed Gull nests have fluctuated between 5 and 18 since 2000. 2022 saw an upturn to 11 nests following a steady decline since 2015, however this declined to 4, the lowest recorded, in 2023. In 2024 there were 4 nests recorded at Ballard Down and one recorded at Durlston.
- 4.20 The boat surveys provide comparable data each year, but is not necessarily a complete count (for example, where not all areas are visible from the sea including overgrown vegetation at Old Harry).
- 4.21 The UK trend shows a decline between 2000 and 2006 but a more recent upturn in 2022 which was reflected in Dorset (see Figure 5), where the decline had been steeper (note that the tiny population size means that small changes result in a large proportional change).




Figure 5: Change numbers of Great Black-backed Gull AONs and the UK index of abundance.

4.22 The 20th century saw both a widespread expansion of the Great Black-backed Gull breeding range and an increase in numbers. The abundance of Great Black-backed Gulls decreased a little between the first census in 1969/70 and 2000. Between 1986 and 2010, abundance peaked in 1999 at 115% of the 1986 reference level but has since decreased by around 20%. It is suggested that Great Black-backed Gulls have a competitive advantage over other scavenging seabirds, such as Fulmar and Herring Gull, and are more adaptable, also taking natural prey (such as Rabbits) and that this may explain why, until recently, they have not undergone the declines experienced by other scavengers.

4.23 Great Black-backed Gull is currently an Amber Listed Bird of Conservation Concern due to a non-breeding population decline (Eaton et al., 2015).

Kittiwake

Following rapid expansion throughout the 1960's and 1970's, the Kittiwake population in Purbeck declined almost as rapidly and in 2024 the only remaining colony (at Blackers Hole) was again small, with just 5 AOS (an increase of 1 from 2023). The future security of a breeding population at this site is therefore extremely uncertain. There is some discussion as to whether the Blackers Hole birds may have moved to Portland Bill, where a colony has re-established itself after a 10-year gap. The decline in Purbeck is steeper than the UK trend.

4.24 Kittiwakes are known to have been present around Durlston in the 1880's (see Lake *et al.* 2011), but only two were recorded by 1957. This site remained the only colony until the late 1960's/early 1970's, when four more sites were colonised and by 1980 the overall population peaked at nearly 300 AONs. After this, all the colonies declined rapidly, and since the mid-1990's, only the Blackers Hole colony has persisted in Purbeck. Despite a brief increase in the mid-2000's, and a slight upturn in 2019 and 2021, the Blackers Hole colony is declining.

4.25 Changes in the Purbeck population mirror the UK trend (see Figure 6), although the population may have peaked earlier, and the decline occurred more rapidly until it slowed in the 21st century. The 2019 slight national upturn was reflected in Purbeck, although numbers dropped again in 2022.

4.26 There used to be a colony of breeding Kittiwakes on Portland Bill (about 33km west), but this similarly declined to the point where breeding ceased around 10 years ago, and birds were no longer seen in the vicinity. However, in 2023 pairs appeared to settle on the cliff ledges below the QinetiQ compound (accurate counts are not possible as there is no access to the area), and this behaviour was repeated in 2024.




Figure 6: Change in numbers of AONs of Kittiwakes in Purbeck and the Purbeck and UK indices of abundance from 1985.

4.27 Nationally, declines in productivity have been related to declines in Sand Eel abundance and are, in some regions, negatively correlated with surface sea temperature (Frederiksen et al., 2004). Kittiwakes are particularly vulnerable to food shortages, as they are surface feeders, and are only able to reach prey on or near the surface. Kittiwakes are Red Listed (Eaton et al., 2015) due to the decline and degree of localisation of the breeding population.

Guillemot

After large declines up to the mid-20th century, Guillemot numbers in Purbeck stabilised in the 1970's and increased overall throughout the 1990's and 2000's, then more rapidly from 2014 onwards. In 2023, the population dropped to 65% of the peak count in 2022. However, in 2024 the population appears to have recovered to a similar level to that seen in 2023, with 1,547 adults counted on ledges. This is the second highest count since the mid-1960's (when systematic recording began). The Purbeck colonies have followed a similar trend to that shown by the UK index of abundance, although fluctuating more widely.

4.28 The Guillemot population is found between Durlston and St. Aldhelm's Head. The number of Guillemots in Purbeck declined from an estimated 2,500-3,500 in the 1930's to about one quarter of this (around 700) in the 1970's (see Lake *et al.* 2011 for more details). After this the overall population began to increase, mainly at the Durlston colony, but also between Crab Rock and Sutton Rock from the early 2000's. In 2024, the colony had mainly increased at Durlston, Blackers Hole, and Sutton Rock.

4.29 Historically there has been variation between the trends at different colonies (see Figure 7). In 2024, there was an increase in all colonies, including the re-emergence of a small number of individuals at White Ware – a nest site not used since the mid-1970's.

4.30 Changes in the Purbeck population correlate broadly with changes in the national index of abundance, although show more fluctuations (see Figure 8).




Figure 7: Changes in numbers of Guillemot individuals at breeding colonies in Purbeck since 1965.




Figure 8: Changes in the total number of Guillemot individuals recorded at breeding ledges in Purbeck compared to the UK index of abundance.

4.31 *The reasons for the national increase are not known, although the recent levelling out may be due to density-dependent effects on breeding success (with competition for space and food becoming critical). Observed low UK productivity, thought to be due to food shortages, combined with low return rates at sampled colonies, suggests that, should productivity decline further, future declines may be likely nationally (JNCC 2011). Guillemot is an Amber Listed Bird of Conservation Concern due to its degree of localisation (Eaton et al. 2015).*

Razorbill

Razorbills declined substantially in Purbeck between 1880 and the early 1960's (when systematic counts began). The overall population continued to decline before stabilising in the 1970's, then fluctuated widely until more steady increases were seen after 2008. By 2022, the number of individual birds (194) was the highest since systematic recording began; although this dropped to 155 in 2023, the population has returned to similar levels in 2024 with a population total of 188 - the second highest count. The population varied notably between colonies. The trend generally reflects that of the UK as a whole, with larger fluctuations greater than those seen in the UK index of abundance.

4.32 Razorbills breed between Durlston and St. Aldhelm's Head in Purbeck. Razorbill was considered to breed on the Purbeck Coast in greater numbers than Guillemot in the 1880's (see Lake *et al.* 2011 for more details). However, by 1932 only 130 birds were recorded, and this total fell further to 58 by 1967 and to just 14 by 1970, by which time many colonies had disappeared altogether. The population then fluctuated but remained steady overall until the late 1980's, after which three crashes, each roughly a decade apart, were followed by recoveries to higher peaks. Substantial increases in 2014, 2018, 2019, and 2022 followed, and although numbers declined by a quarter in 2023, at 188 in 2024 the count is the second highest since systematic recording began in 1964.

4.33 In general, changes have been fairly consistent between the larger colonies (see Figure 9). In 2024, the largest colony was recorded at Crab Hole with 55 individuals, the largest number recorded at this location. Funnel & Reform and Bird Rock saw a decrease on the previous year, whereas the Durlston colony remained stable with a slight increase. No birds were counted at Ragged Rocks and White Ware, or between Hedbury and Halswell.




Figure 9: Changes in counts of individual Razorbills at main colonies between 1965 and 2024 (Durlston birds were not counted in 2020).




Figure 10: Changes in the counts of individual Razorbills and the UK and Purbeck indices of abundance.

4.34 *The Purbeck population has shown large fluctuations since the 1950s (although note that the small size of the population means a small change in numbers results in a large percentage change) (see Figure 10). These fluctuations can obscure overall trends, but since 2008 there has been a clear overall upward trend which is steeper than the fluctuating upward trend in the UK as a whole.*

4.35 *As with Guillemots, it has been suggested that the levelling out seen in the UK index in the 2000's may be due to density dependent mechanisms (JNCC 2011). UK Razorbill productivity has declined steadily since 1993 (possibly due to food shortages), and unless this trend reverses, a continuing overall decline is predicted (JNCC 2011). Razorbill remains an Amber Listed Bird of Conservation Concern due to its degree of localisation (Eaton et al. 2015).*

Puffin

The tiny Puffin population may be stable, although no birds were recorded on the boat survey. The Purbeck population declined severely in the 20th century. By the time the population steadied in the 1990's, the estimated number of breeding pairs was about three and is thought to have fluctuated between one and three since then. In contrast to Purbeck, the national trend was of a significant increase in the last quarter of the 20th century, but more recent monitoring at a small number of large colonies has shown declines in numbers, survival, and productivity.

4.36 Puffins were thought to be abundant in Purbeck at least until 1939 (see Lake et al., 2011) but by 1958 there were only 85 individuals recorded, dropping to 23 in 1975. The population subsequently declined much more slowly until the mid-1990's, after which it stabilised at around two to three breeding pairs (see Figure 11) – a 92% decline since 1958.

4.37 No birds were recorded during the 2023 boat survey. Birds were observed from the cliff and it was estimated that there were a total of 3 breeding pairs from observations of courting behaviour, mating, and birds carrying nesting material (R. Caldow, pers. comm.). However, no birds were observed carrying fish into the nest site and fewer birds were observed during most of June, with an increase in individuals observed towards the end of June / early July. This suggests that breeding was unsuccessful this year. The maximum count of individuals seen this year was nine (R. Caldow, pers. comm.).




Figure 11: Numbers of individual Puffins recorded and estimated number of breeding pairs between 1958 and 2022 (no estimates of pairs were made in 2021 and 2022). Note that the 2023 and 2024 counts were not from the boat survey.

4.38 *The downward trend in Puffin numbers in Purbeck does not reflect the overall increase suggested by UK census returns between 1969 and 2002. However, although UK-wide data are not available for more recent years, monitoring results from two large colonies show subsequent declines. Productivity has fluctuated but appears to have been lower since the 1990s. Caution should be used in drawing wider geographical conclusions from these data. On Lundy Island, where conditions for Puffins have improved through the eradication of rats, numbers have increased from just five individuals in 2006 to over 300 in 2016. Puffins are Amber Listed due to their degree of localisation and categorisation as a species of European Conservation Concern (Eaton et al., 2009).*

References

Eaton, M., Aebischer, N. J., Brown, A., Hearn, R., Lock, L., Musgrove, A., Noble, D., Stroud, D., & Gregory, R. (2015). Birds of Conservation Concern 4: The population status of birds in the UK, Channel Islands and Isle of Man. *British Birds*, 108, 708–746.

Eaton, M. F., Brown, A. F., Noble, D. G., Musgrove, A. J., Aebischer, N. J., Gibbons, D. W., Evans, A., & Gregory, R. D. (2009). Birds of Conservation Concern 3: The population status of birds in the United Kingdom, Channel Islands and the Isle of Man. *British Birds*, 102, 296–341.

Frederiksen, M., Wanless, S., Harris, M. P., Rothery, P., & Wilson, L. J. (2004). The Role of Industrial Fisheries and Oceanographic Change in the Decline of North Sea Black-Legged Kittiwakes. *Journal Of Applied Ecology*, 41, 1129–1139.

Harris, M. P., & Wanless, S. (1996). Differential responses of Guillemot *Uria aalge* and Shag *Phalacrocorax aristotelis* to a late winter wreck. *Bird Study*, 43, 220–230. <https://doi.org/10.1080/00063659609461014>

Haysom, T. S. (1993). The status of some Purbeck sea birds 2. *Dorset Natural History and Archaeological Society*, 144, 215–220.

Haysom, W. T. (1967). A Survey of the Breeding Birds of Purbeck, 88 84-92. *Proceedings of the Dorset Natural History and Archaeological Society*, 88, 84–92.

Haysom, W. T. (1977). The status of some Purbeck seabirds. *Dorset Natural History and Archaeological Society*, 99, 97–103.

JNCC. (2011). *Seabird Population Trends and Causes of Change: 2011 Report Updated April 2011*. Joint Nature Conservation Committee. <http://www.jncc.gov.uk/page-3201>

Lake, S., & Caals, Z. (2022). *Purbeck Seabird Survey 2021* (644). Footprint Ecology /

National Trust.

Lake, S., Liley, D., Lane, K., Hopper, N., & Brereton, T. (2011). *Seabird Breeding Success*

Survey, for Ballard Cliff (Handfast to Ballard Point), Durlston to St. Aldhelms Head and

Gad cliff to White Nothe, South-East Dorset Coast, Dorset. Footprint Ecology /

MARINELife / National Trust.

Lake, S., & Rush, E. (2023). *Purbeck Seabird Survey 2023* (749). Footprint Ecology /

National Trust.

Stanbury, A., Eaton, M., Aebischer, N., Balmer, D., Noble, D., Douse, A., Lindley, P.,

McCulloch, N., & Win, I. (2021). The status of our bird populations: The fifth Birds

of Conservation Concern in the United Kingdom, Channel Islands and Isle of Man

and second IUCN Red List assessment of extinction risk for Great Britain. *British*

Birds, 114(December), 723–747.

Thompson, P. M. (2004). Identifying drivers of change; did fisheries play a role in the

spread of North Atlantic fulmars? In *Management of marine ecosystems:*

Monitoring change in upper trophic levels. Eds I.L. Boyd & S. Wanless. Cambridge

University Press.

Walsh, P. M., Halley, D. J., Harris, M. P., del Nevo, A., Sim, I. M. W., & Tasker, M. L. (1995).

Seabird monitoring handbook for Britain and Ireland: A compilation of methods for

survey and monitoring of breeding seabirds.

http://www.jncc.gov.uk/PDF/pub95_SeabirdHandbook.pdf